

Resonant State in ${}^4_{\Lambda}\text{He}$

D. E. Kahana, S. H. Kahana, and D. J. Millener

Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA

(Dated: February 8, 2008)

In a recent experiment E906 at the BNL-AGS, a search for light $S = -2$ hypernuclei, strong evidence for the nuclide ${}^4_{\Lambda\Lambda}\text{H}$ was found. One of the most striking components of this data was the appearance of a narrow low-momentum π^- line at $k_\pi = 104 - 105$ MeV/c. This was ascribed to the decay of ${}^4_{\Lambda\Lambda}\text{H}$ into a resonant state in ${}^4_{\Lambda}\text{He}$. The existence of such a state is shown to be plausible and its characteristics delineated.

PACS numbers: 21.80+a

Several candidates [1, 2, 3, 4] for doubly-strange hypernuclei have been identified in emulsion experiments. By the nature of the latter approach, usually only one example of a given species is uncovered in each search. The BNL-AGS experiment E906 [5], a counter experiment looking for correlated two π^- decays, was able to accumulate several tens of plausible samples of the hypernucleus ${}^4_{\Lambda\Lambda}\text{H}$. In this work we concentrate on one of the prominent decay modes of this nuclide, ascribed [5, 6] to initial decay into a resonant state in ${}^4_{\Lambda}\text{He}$. This is in analogy to the well-known decay of ${}^5_{\Lambda}\text{He}$ which gives rise to a rather narrow peak in the π^- spectrum [7]. We simply extend the analysis of E906 in a most natural fashion.

One expects the major sequential decays of ${}^4_{\Lambda\Lambda}\text{H}$ to be:

$${}^4_{\Lambda\Lambda}\text{H} \rightarrow {}^4_{\Lambda}\text{He} + \pi^-_H \quad (\sim 112 - 118 \text{ MeV/c}) \quad (1)$$

$${}^4_{\Lambda}\text{He} \rightarrow {}^3\text{H} + p + \pi^-_L \quad (\sim 85 - 95 \text{ MeV/c}) \quad (2)$$

and in particular, a decay into a possible excited state of ${}^4_{\Lambda}\text{He}$,

$${}^4_{\Lambda\Lambda}\text{H} \rightarrow {}^4_{\Lambda}\text{He}^* + \pi^-_L \quad (\sim 104 - 105 \text{ MeV/c}) \quad (3)$$

$${}^4_{\Lambda}\text{He}^* \rightarrow {}^3_{\Lambda}\text{H} + p \quad (4)$$

$${}^3_{\Lambda}\text{H} \rightarrow {}^3\text{He} + \pi^-_H \quad (114.3 \text{ MeV/c}), \quad (5)$$

where π^-_H and π^-_L refer to the high and low momentum members of a correlated pair seen in the experiment.

The resonance depicted in Eqs. (3-4), as indicated below, would certainly be suppressed in the standard single hypernucleus search, for example in the simple ${}^4\text{He}(K^-, \pi^-){}^4_{\Lambda}\text{He}$ reaction [8]. Nevertheless, such a state plays a dominant role in the E906 study, seemingly the only explanation [5] for the strong π^- decay line seen at $104 - 105$ MeV/c. Such a state can be shown to be theoretically plausible, arising naturally in a model [6] consistent with the known ground state of ${}^4_{\Lambda}\text{He}$. The approach followed here is to identify a candidate, not surprisingly a (proton + ${}^3_{\Lambda}\text{H}$) relative p state, and establish its suitability. A state of this architecture would lead directly via the decay of ${}^3_{\Lambda}\text{H}$, to the higher momentum π^- decay line near 114 MeV/c correlated with the considerably narrower feature at $104 - 105$ MeV/c in the E906 data [5].

As was demonstrated in Ref. [9], this state has the characteristics for being a conspicuous daughter for the

decay mode cited above in Eq. (3). As we indicate later, we can construct simple non-spurious $1\hbar\omega$ shell-model states which should be populated strongly in the π^- decay of ${}^4_{\Lambda\Lambda}\text{H}$, and should themselves decay via proton rather than Λ emission. Ideally one might attempt to treat the specific 4-body nature of the ${}^4_{\Lambda\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{He}$ systems [9, 10, 11]. However, practically one is then limited to a restricted selection of possible two-baryon interactions. Even in the $S = -1$ baryon-baryon sector, the known forces, constrained by available data, are by no means uniquely defined, and hence perhaps as yet unreliable. To establish the presence of a weakly bound, or in particular a low-lying resonant, state is then a daunting theoretical task. Indeed, existing many body calculations [12] for the weakly bound ${}^3_{\Lambda}\text{H}$ indicate that $\Lambda\text{N}-\Sigma\text{N}$ coupling must be included, as it must for all the s -shell hypernuclei [13]. This adds to the complexity of such calculations. Ultimately, it is precisely the experimental evidence considered here and elsewhere that should constrain and illuminate hypernuclear few-body theory.

At present, as already indicated, reasonable evidence exists for such a state from experiment [5]. The dynamics of an essentially (p + ${}^3_{\Lambda}\text{H}$) p-state can perhaps be well approximated by a one body + cluster potential problem. The effective interaction between a proton and a three particle hyperon core can be viewed as the average potential arising *after* a combined many-body, configuration mixing, treatment. One is here essentially only interested in the basic s-wave to p-wave separation obtaining, very much an effective single-particle property. That the anticipated resonance emerges naturally from such modeling is suggestive.

The characteristics of such a potential are fairly clear, one is dealing with rather spatially extended systems for both the weakly bound ${}^4_{\Lambda\Lambda}\text{H}$ and resonant ${}^4_{\Lambda}\text{He}^*$. Moreover, the potential can be strongly constrained, by the requirement that its depth also accommodate the ground state of ${}^4_{\Lambda}\text{He}$. The spatial extension suggests one might use a potential with a relatively large radius, or even a surface potential. The resulting spatially large state also favors the decay into the resonance in Eq. (3) from the expected weakly bound ${}^4_{\Lambda\Lambda}\text{H}$.

To help pick out possible candidates for the negative-parity resonance in ${}^4_{\Lambda}\text{He}$, and to illustrate important features of the π^- weak decay of ${}^4_{\Lambda\Lambda}\text{H}$, we enumerate in




FIG. 1: Spectrum and decay thresholds for ${}^4_{\Lambda}\text{He}$. The postulated negative-parity resonant states are indicated above the proton threshold.

Eqs. (6)-(8) the possible $1\hbar\omega$, $T = 1/2$, states in an harmonic oscillator shell-model basis with $\hbar\omega_N = \hbar\omega_{\Lambda}$. Although harmonic oscillator radial wave functions are not appropriate for the loosely bound or resonant states, this basis nevertheless has the virtue that the spurious center of mass states can be eliminated, as in Eq. (6), and the explicit intrinsic-spin structure of the states makes it clear which states can be fed strongly in the π^- weak decay of ${}^4_{\Lambda}\text{H}$.

$$\sqrt{\frac{\mu}{3+\mu}} (s^2p)[3]1/2 \times s_{\Lambda} - \sqrt{\frac{3}{3+\mu}} s^3 \times p_{\Lambda} \quad (6)$$

$$(s^2p)[21]1/2 \times s_{\Lambda} \quad (7)$$

$$(s^2p)[21]3/2 \times s_{\Lambda} \quad (8)$$

Here $\mu = m_{\Lambda}/m_N$ and $[f]S_{3N}$ labels the spatial symmetry and intrinsic spin of the three nucleons. For each of the three classes of states $L = 1$ and two values of total intrinsic spin S are possible. In the same model, the ${}^4_{\Lambda}\text{H}$ initial state is simply $s^2(1) \times s^2_{\Lambda}(0)$ with $S = 1$. Because the dominant piece of the weak-decay operator does not involve intrinsic spin, we need consider only ${}^4_{\Lambda}\text{He}$ states with $L = 1$, $S = 1$, and $J = 0, 1, 2$.

Because the proton threshold in ${}^4_{\Lambda}\text{He}$ at 7.75 MeV is much higher than the Λ threshold at 2.39 MeV (see Fig. 1) any state with even a small p_{Λ} component will decay to ${}^3\text{He} + \Lambda$. This rules out the states of Eq. (6), although such states could be fed via the s_{Λ} component in the weak decay leading to a ${}^3\text{He} + \Lambda$ final state. The states in Eqs. (7) and (8), which should be at similar excitation energies, qualify as candidates. We do not expect the relatively weak ΛN interaction to mix these configurations strongly with those of Eq. (6). We note that in light nuclei a number of narrow states exist at high

TABLE I: Matrix elements for the production of single- Λ hypernuclear configurations in the π^- weak decay of ${}^4_{\Lambda}\text{H}$ are given in the first row in units of $s_{\pi^-}^2 \langle l_N | j_{l_N}(k_{\pi}r) | s_{\Lambda} \rangle^2 / 4\pi$ where $\mu/(3+\mu) = 0.284$. The remaining rows specify the breakup of the single- Λ hypernuclear configurations under the assumptions described in the text. Combining production and decay in this simple approach shows that ${}^3_{\Lambda}\text{H} + p$ is favored over $d + p + \Lambda$ by a factor of 11/3 to 4/3.

Final state	${}^4_{\Lambda}\text{He}(1^+)$	Eq. (6)	Eq. (7)	Eq. (8)
Production	1	$\mu/(3+\mu)$	1	4
${}^4_{\Lambda}\text{He}(1^+)$	1			
${}^3\text{He} + \Lambda$		1		
${}^3_{\Lambda}\text{H} + p$			1/9	8/9
$d + p + \Lambda$			8/9	1/9

excitation energies because of the symmetry structure of the wave functions, e.g. a $3/2^+$ state of $t + d$ structure at 16.75 MeV, in ${}^5\text{He}$ and an analogous state in ${}^5\text{Li}$ despite a large decay energy into the $\alpha + N$ channel. If the intrinsic spins in the configurations in Eqs. (7) and (8) are recoupled to the form $(s^2s_{\Lambda})S_{3H} \times p$, $S_{3H} = 1/2$ is required for the ${}^3_{\Lambda}\text{H} + p$ channel; $S_{3H} = 3/2$ leads to the $d + \Lambda + p$ final state. In fact, the configuration in Eq. (8) dominates the $S_{3H} = 1/2$ strength. The preceding discussion relating to the possible decay channels of single- Λ hypernuclear configurations produced in the π^- weak decay of ${}^4_{\Lambda}\text{H}$ is summarized in rows 2–5 of Table I.

The weak-decay of ${}^4_{\Lambda}\text{H}$ has been treated by Kumagai-Fuse and Okabe [9]. The widths that they calculate for decays to ${}^4_{\Lambda}\text{He}(1^+)$ and to ${}^3_{\Lambda}\text{H} + p$ are $0.68\Gamma_{\Lambda}$ and $0.80\Gamma_{\Lambda}$, respectively. We reproduce the essential features of these results using just the dominant piece of the weak-decay operator which does not involve the nucleon spin. In the plane-wave limit for the outgoing π^- the operator is simply $s_{\pi^-} j_{\lambda}(k_{\pi}r) Y_{\lambda}$, where s_{π^-} is a strength known from the decay of the free Λ [9]. The squares of the matrix elements of this operator taking into account the sum over J_f with the usual statistical weighting of $(2J_f + 1)/(2J_i + 1)$ (Eqs. (9) and (10) of [9]) are given in the first row of Table I. Putting in the kinematic factors (and the small contribution from the spin-dependent weak-decay operator with a strength given by p_{π^-} [9]) gives a width $0.806 \langle s_N | j_0(k_{\pi}r) | s_{\Lambda} \rangle^2 \Gamma_{\Lambda}$ to the ${}^4_{\Lambda}\text{He}(1^+)$ state, where the radial matrix element is not very much smaller than unity because $k_{\pi} \sim 0.53 \text{ fm}^{-1}$ is quite small. The radial matrix element for the decays leading to ${}^3_{\Lambda}\text{H} + p$ is smaller and a rough estimate suggests that it compensates for the 11/3 factor for production noted in the caption of Table I to give a rate comparable to that for the ${}^4_{\Lambda}\text{He}(1^+)$ state. The point is that there is quite good agreement with Ref. [9] and that ${}^3_{\Lambda}\text{H}$ should be produced sufficiently that the scenario [5] put forward in Eqs. (3), (4), and (5) is reasonable.

The next order of business is to construct an average potential describing a proton in both the ground and

resonant states of ${}^4_\Lambda\text{He}$. We reemphasize that we are using such a potential to represent the average interaction between the proton and an effective ${}^3_\Lambda\text{H}$ cluster. We employ a resonance code due to T. Vertse, K. F. Pal and Z. Balogh [14] to perform the necessary calculations, together with a surface Saxon-Woods potential. The addition of a spin-orbit potential doesn't change the basic results and, in any case, the configurations in Eqs. (7) and (8) correspond to mixtures of $p_{1/2}$ and $p_{3/2}$ proton orbits. Thus, we take,

$$V^{Surf}(r) = -V_0 \left[\frac{4e^{(r-R)/a}}{(1 + e^{(r-R)/a})^2} \right], \quad (9)$$

with $R = r_0 A^{1/3}$ the potential radius and a the diffusivity. A Coulomb potential, important to constraining the width of a resonant proton line, is included. This potential is taken as that of a uniform charge distribution with radius parameter $r_{0c} = r_0$.

The ground state proton has of course $l = 0$ and a known binding. Only the strength of the surface potential component is varied and reproduces the correct 7.75 MeV separation energy with a depth $V_0 = 28.09$ MeV, radius parameter $r_0 = 1.40$ fm, and diffusivity $a = 0.5$ fm. It should be noted that the actual nuclear radius is $R = 2.07$ fm since the A used in the resonance code [14] is the hypernuclear core mass 3.21 amu. In this completely specified well, the p -wave resonance appears at $\epsilon = 1.18$ MeV with a width of 1.00 MeV. Both these numbers are consistent with the observed narrow π^- line at 104 – 106 MeV/c in E906, recalling that the resolution in this experiment was ~ 2.5 MeV. For the choice $r_0 = 1.45$ fm, the corresponding well and resonance parameters are $V_0 = 27.68$, $\epsilon = 0.97$ MeV, and width 0.70 MeV.

Another possible variation is to fix the proton separation energy at 7.18 MeV, from averaging the ${}^4_\Lambda\text{He}$ ($0^+, 1^+$) ground state doublet positions. Resetting the radius parameters to $r_0 = 1.45$ fm, the p -wave resonance appears at 1.19 MeV with a width of 1.04 MeV in a slightly shallower surface well with $V_0 = 26.69$ MeV. One is, of course, comparing only to centroids of the experimental features.

Most convincing in this analysis is the ease with which the ${}^4_\Lambda\text{He}$ resonance is extracted, once the ground state binding of the proton is assigned. The final resonance parameters obtained cannot be taken too literally, but the energy and width have little variability with reasonable changes in other potential parameters.

With the resonance energy calculated the $\Lambda\Lambda$ pairing energy $\Delta B_{\Lambda\Lambda}$ can be estimated from the position of the narrow π^- peak ascribed to the weak decay from ${}^4_\Lambda\text{H}$ to ${}^4_\Lambda\text{He}^*$. The π^- momentum k^* is given by

$$\epsilon_\pi + k^{*2}/2M({}^4_\Lambda\text{He}^*) = M({}_\Lambda\Lambda^4\text{H}) - M({}_\Lambda\Lambda^4\text{He}^*). \quad (10)$$

To a good approximation in the region of momentum considered here,

$$k_\pi^* = (107.466 - 1.6391\Delta) \text{ MeV/c}, \quad (11)$$

where $\Delta = B^* + \epsilon_R$ and $B^* = 2\bar{B}_\Lambda({}^3_\Lambda\text{H}) + \Delta B_{\Lambda\Lambda}$ is the full binding energy of the Λ pair in ${}^4_\Lambda\text{H}$ excluding rearrangement effects and coupling to other hyperon channels. These relations exhibit the dependence of the measured meson momentum on the combined resonance energy and $\Lambda\Lambda$ interaction energy. In evaluating the contribution of the ΛN interaction, a recoupling of the initial $s^2 \times s^2_\Lambda$ wavefunction into $(s^2 s_\Lambda)S_{3H}$ coupled to a spectator s_Λ leads to a statistical weighting of B_Λ values for the real $1/2^+$ hypertriton and the unbound and unknown $3/2^+$ hypertriton denoted by $\bar{B}_\Lambda$. Using the known value for $2B_\Lambda({}^3_\Lambda\text{H})$ in place of $2\bar{B}_\Lambda$ as we do below means that $\Delta B_{\Lambda\Lambda}$ will be underestimated (but not by a large amount). In our model, for example, taking $r_0 = 1.40$ fm, hence $\epsilon_R = 1.18$ MeV and $\Delta B_{\Lambda\Lambda} = 0.34$ MeV, puts the centroid of the decay momentum at $k^* = 104.5$ MeV/c, nominally the position of the measured centroid for this decay momentum. Alternatively, $\epsilon_R = 0.97$ MeV and the same decay momentum results in $\Delta B_{\Lambda\Lambda} = 0.55$ MeV.

The indefiniteness in the measured π_L momentum [5] subjects the pairing energy to some uncertainty. For example, with $\epsilon_R = 1.18$, lowering k^* by 300 keV results in $\Delta B_{\Lambda\Lambda} = 0.55$ MeV, while a corresponding increase in k^* yields 0.17 MeV. Whatever the choice, $\Delta B_{\Lambda\Lambda}$ here remains small and appreciably less than the ~ 1 MeV determined for ${}^6_\Lambda\text{He}$ [4].

One very interesting feature of these estimated values for $\Delta B_{\Lambda\Lambda}$, say 0.34 MeV, is the momentum for the truly two body decay

$${}^4_\Lambda\text{H} \rightarrow {}^4_\Lambda\text{He}(1^+) + \pi_H^-. \quad (12)$$

A similar approximation to that in Eq. (11) produces, with no ϵ_R involvement,

$$k_\pi(1^+) = (117.454 - 1.5531B^*) \text{ MeV/c}, \quad (13)$$

and $k_\pi = 116.5$ MeV/c. Such a line, although still consistent with the observed broad peak near 115 MeV/c, should be relatively easy to separate, in an improved resolution experiment, from the known 114.3 MeV/c for the decay

$${}^3_\Lambda\text{H} \rightarrow {}^3_\Lambda\text{He} + \pi_H^-. \quad (14)$$

Finding this peak in the π_H^- spectrum would provide more transparent evidence for the existence of ${}^4_\Lambda\text{H}$.

In conclusion, we see that the existence of states above threshold in ${}^3_\Lambda\text{H} + \text{proton}$ system can straightforwardly describe the narrow, low momentum, π^- feature at 104 – 105 MeV/c observed in the BNL experiment E906. An argument in favor of such resonances is that a common average particle-cluster potential can be used to describe the gross structure of both the ground-state doublet of ${}^4_\Lambda\text{He}$ and the negative-parity resonances. To pursue the problem further, one must surely involve the full interaction between the “resonant” proton and its hypernuclear core, including the NN, both the odd and even state ΛN , forces and very likely ΛN - ΣN coupling.

In Ref. [9], a p-wave resonance could not be produced by varying only the poorly known, and probably relatively weak, odd-state ΛN strength. These very light hypernuclei certainly warrant more theoretical study.

The inferred energy of the resonance and energy range possible for $\Delta B_{\Lambda\Lambda}$, the latter likely somewhat less than 0.5 MeV, are not unreasonable. Takahashi *et al.* [4] found $\Delta B_{\Lambda\Lambda} = 1.0 \pm 0.38$ MeV for ${}^6_{\Lambda\Lambda}\text{He}$. One certainly expects a smaller value for the more extended mass 4 system, but the suggested values and their likely errors allow for consistency. Only future experiments with better statistics and better resolution can settle this issue. We reemphasize that, although the analysis followed here is the most natural interpretation of the existing data [5], this interpretation would be much illuminated by such followup experiments.

For the meantime, it is also of high importance to perform many-body calculations [9, 10, 11] with forces constrained by as much data as is available, including of course that from E906. The latter experiment has pointed to the existence of two interesting nuclides, both mass 4 objects, one a very light $S = -2$ hypernucleus, the other an unusual, if not completely unexpected, resonance in an $S = -1$ daughter nucleus.

The initial discoveries of $S = -2$ nuclei [1, 2, 3] had one

measurement in common, the ~ 4.5 MeV value for $\Delta B_{\Lambda\Lambda}$, irrespective of species. This seemed excessively large, considering the expectation of a rather weak $\Lambda\Lambda$ interaction, and hinted at possibly interesting short range behaviour, perhaps even bag-like structure, in these doubly-strange hypernuclei. The value implied by the present analysis and by the recent KEK emulsion experiment [4], $\Delta B_{\Lambda\Lambda} \leq 1.4$ MeV, apparently put paid to such speculation. However, the possibility of unusual short range structure cannot be completely ruled out, and would certainly complicate the calculation of light systems containing a Λ pair. Incidentally, a judiciously selected excited state inserted into the early analyses [1, 3, 15] would also lower the pairing energies extracted there, in line with the recent measurements and the present analysis.

One can also infer, from the small $\Delta B_{\Lambda\Lambda}$ suggested here for ${}^4_{\Lambda\Lambda}\text{H}$, a possible binding of less than 1.0 MeV for the highly elusive H -dibaryon [16]. Production of this object is severely reduced by the strong likelihood of a repulsive core in the $\Lambda\Lambda$ interaction [17]. Evidently, the threshold for study of $S = -2$ nuclei has just been breached; many, and more extensive, searches are needed.

This manuscript has been authored under the US DOE grant NO. DE-AC02-98CH10886.

-
- [1] M. Danysz *et al.*, Nucl. Phys. **49**, 121 (1963); R. H. Dalitz *et al.* Proc. Roy. Soc. Lond. **426**, 1 (1989).
 - [2] D. Prowse, Phys. Rev. Lett. **17**, 782 (1966).
 - [3] S. Aoki *et al.*, Phys. Rev. Lett. **65**, 1729 (1990); S. Aoki *et al.*, Prog. Theor. Phys. **85**, 1287 (1991).
 - [4] H. Takahashi *et al.*, Phys. Rev. Lett. **87**, 212502 (2001).
 - [5] J. K. Ahn *et al.*, Phys. Rev. Lett. **87**, 132504 (2001).
 - [6] D. E. Kahana, S. H. Kahana, and D. J. Millener, Resonances in the Production of $S=-2$ hypernuclei, Poster Session, INPC2001, Berkeley CA, July 2001.
 - [7] W. Gajewski *et al.*, Nucl. Phys. **B14**, 11 (1969); T. Motoba, H. Bandō, T. Fukuda, and J. Žofka, Nucl. Phys. **A534**, 597 (1991); I. Kumagai-Fuse, S. Okabe, and Y. Akaishi, Phys. Rev. C **54**, 2843 (1996).
 - [8] M. J. Athanas, PhD. Dissertation, Carnegie Mellon University, 1992, unpublished; H. Tamura *et al.*, Prog. Theor. Phys. Suppl. **113**, 1 (1994); H. Outa *et al.*, Nucl. Phys. **A639**, 251c (1998).
 - [9] I. Kumagai-Fuse and S. Okabe, Phys. Rev. C **66**, 014003 (2002).
 - [10] I. N. Filikhin, and A. Gal, Phys. Rev. Lett. **89**, 172502 (2002); Nucl. Phys. **A707**, 491 (2002).
 - [11] H. Nemura, Y. Akaishi, and K. S. Myint, nucl-th/0211082.
 - [12] K. Miyagawa and W. Glöckle, Phys. Rev. C **48**, 2576 (1993); K. Miyagawa, *et al.*, Phys. Rev. C **51**, 2905 (1995).
 - [13] Y. Akaishi, T. Harada, S. Shinmura, and K. S. Myint, Phys. Rev. Lett. **84**, 3539 (2000); E. Hiyama *et al.*, Phys. Rev. C **65**, 011301(R) (2001); A. Nogga, H. Kamada, and W. Glöckle, Phys. Rev. Lett. **88**, 172501 (2002); H. Nemura, Y. Akaishi, and Y. Suzuki, Phys. Rev. Lett. **89**, 142504 (2002).
 - [14] T. Vertse, K. F. Pal and Z. Balogh, Comp. Phys. Comm. **27**, 309 (1982).
 - [15] C. B. Dover, D. J. Millener, A. Gal and D. H. Davis, Phys. Rev. C **44**, 1905 (1991).
 - [16] R. L. Jaffe, Phys. Rev. Lett. **38**, 195 (1977); **38**, 617(E) (1977).
 - [17] D. E. Kahana and S. H. Kahana, Phys. Rev. C **60**, 0625206 (2000).